

Dr. Mark Sywak FRACS

Tel: (02) 9437 1731

Mob: 0400 209 039

Email: marksywak@nebsc.com.au

Website: www.nebsc.com.au

Dr. Mark Sywak FRACS

*Suite 202, AMA House
69 Christie Street, St. Leonards 2065*

North Shore Private Hospital
Mater Private Hospital
Manly District Hospital
Royal North Shore Hospital

N O R T H E R N
E N D O C R I N E A N D
B R E A S T S U R G E R Y
C E N T R E

Dr. Mark Sywak FRACS

POST-OPERATIVE
INSTRUCTIONS
MINIMALLY INVASIVE
THYROID SURGERY

*Suite 202, 69 Christie Street
St. Leonards 2065*

*Suite 1, Level 2, 27 Belgrave Street
Manly 2095*

*75 Market Street
Wollongong 2500*



Tel: (02) 9437 1731

Mob: 0400 209 039

Email: marksywak@nebsc.com.au

Website: www.nebsc.com.au

Minimally Invasive Thyroid Surgery

Wound Care: Your wound will be covered with tape, which should be left in place for about 2 weeks. The tape will be removed at your first postoperative visit. You may notice some dried blood under the tape but that is of no concern. You can wash and shower with the tape in place and even get it wet. Once removed or fallen off, the tape may need to be replaced and you will need to change it every 2 or 3 days for the next 2 to 3 months in order to get the best possible cosmetic result. The preferred tape is narrow, flesh coloured Micropore which can be obtained from your chemist.

Activities: You should generally restrict vigorous activities for 1 to 2 weeks after surgery. Activities which involve turning the head suddenly, such as driving in heavy traffic, should be avoided for 5-7 days post surgery, although local driving is acceptable. Commonsense is the best way to avoid straining your neck.

Local symptoms: A variety of local symptoms are common for several weeks after surgery including tightness, swelling of the local area around the wound, choking or having to force down food. Weakness of the voice with prolonged use may also occur. Occasionally a symptom may last for months.

Late Complications: The only delayed complication of concern is wound infection. This may have occurred if the wound becomes very red, hot and more swollen. If that occurs you must seek attention from your local doctor straight away who will arrange for you to have antibiotics.

Follow-up: Generally your follow-up will consist of:

- 1 A visit at 2 to 4 weeks after surgery to have the tape removed and pathology discussed. You will need to ring the office to make an appointment for that visit.
- 2 A second follow-up consult at 3 months.

In addition, if you have been placed on calcium supplements, you need to see by your local doctor on a weekly basis to have a blood test to check your calcium level, and to have your calcium supplements progressively reduced (see "Calcium Supplements" below). You will also need to be reviewed by your local doctor or endocrinologist at around 6 weeks after surgery and have thyroid function tests to check the function of your remaining thyroid. Your local doctor or endocrinologist may need to see you more frequently for specific review if there are any problems.

Country Patients: If you are unable to return to Sydney for follow-up, it may be possible for the tape to be changed and for your follow-up to be undertaken by your local doctor. We would need to confirm these arrangements before you leave the hospital.

Calcium Supplements: A temporary drop in calcium levels may sometimes occur even after minimal access thyroid surgery. The calcium level is checked in hospital and if it is abnormal, you will be sent home on calcium supplements (Caltrate). You will then need to see your local doctor every week after discharge to have a blood test and to have the calcium levels checked. If, at each visit, the calcium level in the blood is normal, then the dose needs to be reduced according to the protocol below. Please give this protocol to your local doctor.

If you have any problems, following your thyroid surgery, you should contact Dr. Sywak or your local doctor.

IF YOU HAVE BEEN DISCHARGED ON CALCIUM MEDICATION, PLEASE DETACH THIS PAGE AND GIVE IT TO YOUR DOCTOR ONE WEEK AFTER DISCHARGE

Dear Doctor:

Your patient has been discharged on calcium supplements following their thyroid surgery. They have been asked to see you on a weekly basis to have their serum calcium level checked and their medication reduced according to the protocol:

On discharge they will normally be on:

2 Caltrate twice a day

If calcium is normal at one week reduce to:

1 Caltrate twice a day

If calcium is normal the next week reduce to:

1 Caltrate daily

If calcium is normal the next week:

Cease Caltrate

DOCTOR, IF YOU HAVE ANY QUESTIONS OR CONCERNS PLEASE CONTACT MARK SYWAK AS LISTED ON THE OVER LEAF

Dr. Mark Sywak FRACS