

**Clinical Associate Professor
Mark Sywak FRACS**

**POST-OPERATIVE
INSTRUCTIONS:
HEMITHYROIDECTOMY
& PARTIAL
THYROIDECTOMY**



AMA HOUSE

*Level 2, Suite 202, 69 Christie St,
St. Leonards 2065*

NORTHERN BEACHES HOSPITAL

*Suite 9, Level 6, 105 Frenchs Forest
Road, Frenchs Forest 2086*

**Clinical Associate Professor
Mark Sywak FRACS**

Tel: (02) 9030 1652

Mob: 0400 209 039

Email: marksywak@nebsc.com.au

Website:

www.SydneyEndocrineSurgery.com.au

**Clinical Associate Professor
Mark Sywak FRACS**

North Shore Private Hospital

Mater Private Hospital

Northern Beaches Hospital

Royal North Shore Hospital

Tel: (02) 9030 1652

Mob: 0400 209 039

Email: marksywak@nebsc.com.au

Website:

www.SydneyEndocrineSurgery.com.au

Hemithyroidectomy & Partial Thyroidectomy

Wound Care: Your wound will be covered with tape, which should be left in place for about 1-2 weeks. The tape will be removed at your first postoperative visit. You may notice some dried blood under the tape but that is of no concern. You can wash and shower with the tape in place and even get it wet. Once removed or fallen off, the tape needs to be replaced and you will need to change it every 2 or 3 days for the next 2 to 3 months in order to get the best possible cosmetic result. The preferred tape is narrow, flesh coloured Micropore which can be obtained from your chemist.

Activities: You should generally restrict vigorous activities for 1 to 2 weeks after surgery. Activities which involve turning the head suddenly, such as driving, lifting over 10kg from the ground up and gym workouts.

Local symptoms: A variety of local symptoms are common for several weeks after surgery including tightness, choking or having to force down food. Headaches and tiredness are also common as is weakness of the voice with prolonged use. Swelling of the neck around the wound is also common and may benefit from daily massage with Bio Oil or Vitamin E cream but not to be used within the first 8 weeks post-surgery. Numbness of the skin above the wound may be present and may last for many months.

Late Complications: The only delayed complication of concern is wound infection. This may have occurred if the wound becomes very red, hot and more swollen. If that occurs you must seek attention from your local doctor straight away who will arrange for you to have antibiotics.

Follow-up: Generally your follow-up will consist of:

- 1 A visit at 1 to 2 weeks after surgery to have the tape removed & pathology discussed. If you do not have a post-operative appointment you will need to ring the office to make an appointment.
- 2 A second follow-up consult at 3 months if needed.

Country Patients: If you are unable to return to Sydney for follow-up, it may be possible for the tape to be changed and for your follow-up to be undertaken by your local doctor. We would need to confirm these arrangements before you leave the hospital.

Calcium Supplements: A temporary drop in calcium levels may sometimes occur even after hemithyroidectomy or partial thyroidectomy. The calcium level is checked in hospital and if it is abnormal, you will be sent home on calcium supplements (Caltrate). You will then need to see your local doctor every week after discharge to have a blood test and to have the calcium levels checked. If, at each visit, the calcium level in the blood is normal, then the dose needs to be reduced according to the protocol below. Please give this protocol to your local doctor.

If you have any problems, following your thyroid surgery, you should contact A/Prof Sywak or your local doctor.

IF YOU HAVE BEEN DISCHARGED ON CALCIUM MEDICATION, PLEASE DETACH THIS PAGE AND GIVE IT TO YOUR DOCTOR ONE WEEK AFTER DISCHARGE

DR SYWAK WILL ADJUST YOUR CALCIUM MEDICATION ON THE FIRST POST-OPERATIVE VISIT.

IF YOU ARE FROM OUT OF SYDNEY YOU MAY NEED TO VISIT YOUR GP FOR MONITORING . PLEASE GIVE THIS SECTION TO YOUR GP

Dear Doctor:

Your patient has been discharged on calcium supplements following their thyroid surgery. They have been asked to see you on a weekly basis to have their serum calcium level checked and their medication reduced according to the protocol:

On discharge they will normally be on:

2 Caltrate twice a day

If calcium is normal at one week reduce to:

1 Caltrate twice a day

If calcium is normal the next week reduce to:

1 Caltrate daily

If calcium is normal the next week:

Cease Caltrate

DOCTOR, IF YOU HAVE ANY QUESTIONS OR CONCERNS PLEASE CONTACT

DR. MARK SYWAK AS LISTED ON THE OVER LEAF

**Clinical Associate Professor
Mark Sywak FRACS**